

Edinburgh-/92-93/03

hep-th/9312031

R -matrices for the $n = 2, 3$ Elliptic Calogero-Moser Models

H. W. Braden & T. Suzuki

Department of Mathematics and Statistics,
University of Edinburgh,
Edinburgh, UK.

February 7, 2008

Abstract

The classical (dynamical) R -matrices for the 2- and 3-body Calogero-Moser models with elliptic potentials are given. The 3-body case has an interesting nontrivial structure that goes beyond the known ansatz for momentum independent R -matrices. The R -matrices presented include the dynamical R -matrices of Avan and Talon as degenerate cases of the elliptic potential.

The elliptic Calogero-Moser model [1, 2] is a completely integrable system of n -particles on the line interacting via pairwise (so called type IV) potentials $v(q_{ij}) = a\wp(aq_{ij})$ with $q_{ij} = q_i - q_j$ [3]. The function $\wp$ is the Weierstrass elliptic function, an even doubly-periodic function with periods $2\omega_1$ and $2\omega_2$; the various degenerations of this function also give completely integrable systems. In the limit in which one of the periods goes to infinity the potential reduces to either the (type II) potential $v(q) = a^2/\sinh(aq)^2$ or the (type III) potential $v(q) = a^2/\sin(aq)^2$. When both periods go to infinity we obtain the (type I) potential $v(q) = q^{-2}$. Although the Calogero-Moser model can be generalised to arbitrary semisimple Lie algebras our attention in this letter will be on the original Calogero-Moser model which corresponds to the algebra gl_n .

The model has a Lax pair formulation $i\dot{L} = [L, M]$. The L operator may be expressed as

$$L \equiv \sum_{\mu} L^{\mu} X_{\mu} = \sum_{i=1}^n p_i H_i + i \sum_{\alpha \in \Phi} w_{\alpha} E_{\alpha}. \quad (1)$$

Here $X_{\mu} = \{H_i, E_{\alpha}\}$ is a Cartan-Weyl basis for the algebra gl_n with $\{H_i\}$ a basis for the Cartan subalgebra and $\{E_{\alpha}\}$ the corresponding set of step operators. For the Lie algebra gl_n the root system, which labels the step operators, may be written as $\Phi = \{e_i - e_j, 1 \leq i \neq j \leq n\}$, with e_i being an orthonormal basis of $\mathbb{R}^n$. We may view the positions $(q_1, q_2, \dots, q_n)$ as components of a vector q in this vector space and so for $\alpha = e_i - e_j$ we have $q_{ij} = \alpha \cdot q$. It will be convenient to use the shorthand f_{α} for the function that takes the value $f(\alpha \cdot q)$ when evaluated at q . The function w_{α} in (1) is determined by the requirement that the Lax equation reproduces the Hamiltonian equations of motion. This means the function w_{α} should satisfy the following addition formula,

$$w_{\alpha} w'_{\beta} - w_{\beta} w'_{\alpha} = (z_{\alpha} - z_{\beta}) w_{\alpha+\beta}, \quad \text{where} \quad z_{\alpha}(u) = \frac{w''_{\alpha}(u)}{2w_{\alpha}(u)}. \quad (2)$$

The solutions to this functional equation are given by

$$w_{\alpha} = \frac{\sigma(u - \alpha \cdot q)}{\sigma(u) \sigma(\alpha \cdot q)} e^{\zeta(u) \alpha \cdot q}. \quad (3)$$

Here $\sigma(x)$ and $\zeta(x) = \sigma'(x)/\sigma(x)$ are Weierstrass's sigma and zeta function. The function w_{α} given in (3) depends not only on the coordinate differences $q_{ij} = \alpha \cdot q$ but also on a spectral parameter u [4]. If the spectral parameter u is equal to one of ω_1, ω_2 or $-\omega_1 - \omega_2$ then w_{α} becomes $1/\text{sn } \alpha$, $\text{cn } \alpha/\text{sn } \alpha$ or $\text{dn } \alpha/\text{sn } \alpha$, where $\text{sn } \alpha, \text{cn } \alpha$ and $\text{dn } \alpha$ are the Jacobi elliptic functions. In such cases w_{α} is an odd function, *i.e.*, $w_{-\alpha} = -w_{\alpha}$, and the Lax operator L is hermitian.

The Lax pair formulation immediately shows the functionally independent quantities $I_m \equiv \text{Tr } L^m$ ($m \leq n$) to be conserved. Their Poisson commutativity, $\{I_m, I_k\} = 0$, and hence the complete integrability of the model, follows from the existence of a $\mathfrak{g} \otimes \mathfrak{g}$ -valued R -matix [5, 6, 7], $R = R^{\mu\nu} X_{\mu} \otimes X_{\nu}$, satisfying,

$$\{L \otimes L\} = [R, L \otimes 1] - [R^{\pi}, 1 \otimes L]. \quad (4)$$

Here $R^{\pi} \equiv PRP^{-1}$ with P being the permutation map: $P(a \otimes b)P^{-1} = b \otimes a$. Proving the Poisson commutativity $\{I_m, I_k\} = 0$, or equivalently finding such an R -matrix, is one

of the central tasks in proving the integrability of a model. Recently R -matrices for the Calogero-Moser models with types I, II, III, and V potentials subject to $w_{-\alpha} = -w_{\alpha}$ were constructed by Avan and Talon [8], while spectral parameter dependent R -matrices for the type IV potential were independently given by Sklyanin [9] and the authors [10]. Moreover, we also proved [10, Theorem 4.6] that when $n \geq 4$ no momentum independent R -matrices exist for nondegenerate type IV potentials satisfying $w_{-\alpha} = -w_{\alpha}$. However, as mentioned in [10], the Lie algebra consistency equations that forbid momentum independent R -matrices for $n \geq 4$ are trivially satisfied for $n = 2, 3$. In this letter we will present explicit R -matrices for the $n = 2, 3$ systems with potential satisfying $w_{-\alpha} = -w_{\alpha}$. The structure found in the $n = 3$ case is rather interesting and provides an example of an R -matrix that goes beyond the known ansatz.

Let us start the calculation of the R -matrix. To begin with, we decompose eq.(4) into components with respect to the basis $X_{\mu} \otimes X_{\nu}$. Upon substituting L into (4) and comparing both sides, we obtain three equations,

$$0 = \sum_{\alpha} (R^{\alpha j} \alpha_i - R^{\alpha i} \alpha_j) w_{-\alpha}, \quad (5)$$

$$-\alpha_i w'_{\alpha} = i\alpha \cdot p R^{\alpha i} + \alpha \cdot R^i w_{\alpha} + \sum_{\beta} (\beta_i w_{\beta} R^{-\beta \alpha} + w_{\alpha-\beta} R^{\beta i} c_{\beta \alpha-\beta}^{\alpha}), \quad (6)$$

$$0 = \alpha \cdot R^{\beta} w_{\alpha} - \beta \cdot R^{\alpha} w_{\beta} + i(\alpha \cdot p R^{\alpha \beta} - \beta \cdot p R^{\beta \alpha}) + \sum_{\gamma \in \Phi} (R^{\gamma \beta} c_{\gamma \alpha-\gamma}^{\alpha} w_{\alpha-\gamma} - R^{\gamma \alpha} c_{\gamma \beta-\gamma}^{\beta} w_{\beta-\gamma}). \quad (7)$$

The structure constant $c_{\mu\nu}^{\lambda}$ are defined by $[X_{\mu}, X_{\nu}] = c_{\mu\nu}^{\lambda} X_{\lambda}$ and we have introduced the shorthand notation $\alpha \cdot R^{\beta} = R^{i\beta} - R^{j\beta}$ for $\alpha = e_i - e_j$. The assumption that the R -matrix is momentum independent and that $w_{-\alpha} = -w_{\alpha}$ greatly simplify these equations. We find

$$R^{\alpha i} = 0, \quad R^{\alpha \beta} = 0 \quad \text{if} \quad \alpha \neq \pm \beta, \quad (8)$$

$$R^{\alpha \alpha} + R^{-\alpha - \alpha} = 0, \quad R^{\alpha - \alpha} + R^{-\alpha \alpha} = 0, \quad R^{\alpha \alpha} + R^{-\alpha \alpha} = -\frac{w'_{\alpha}}{w_{\alpha}}, \quad (9)$$

and

$$\alpha \cdot R^{\beta} w_{\alpha} - \beta \cdot R^{\alpha} w_{\beta} = c_{\alpha\gamma}^{\beta} (R^{\alpha \alpha} + R^{\beta \beta}) w_{\gamma} + c_{-\alpha\gamma'}^{\beta} (R^{-\alpha \alpha} + R^{-\beta \beta}) w_{\gamma'}. \quad (10)$$

Also $R^{ij} = P^{ij}$, where P^{ij} is a matrix such that $\alpha \cdot P^j = 0 \quad \forall \alpha \in \Phi$ [10, Lemma 4.1]. Note that at most one term on the righthand side of (10) is nonzero. If, for example, $\gamma = \beta - \alpha \in \Phi$ we have only the first term nonvanishing and so $\alpha \cdot R^{\beta} w_{\alpha} - \beta \cdot R^{\alpha} w_{\beta} = c_{\alpha\gamma}^{\beta} (R^{\alpha \alpha} + R^{\beta \beta}) w_{\gamma}$. However (as $-\beta, -\gamma \in \Phi$) we also know that $-\beta = -\alpha - \gamma \in \Phi$ and so we obtain from the second term (after utilizing $c_{-\alpha-\beta}^{-\gamma} = -c_{\alpha\beta}^{\gamma}$) that $\alpha \cdot R^{-\beta} w_{\alpha} - \beta \cdot R^{\alpha} w_{\beta} = c_{\alpha\gamma}^{\beta} (R^{-\alpha \alpha} + R^{\beta - \beta}) w_{\gamma}$. These exemplify some of the consistency requirements mentioned earlier that forbid (nodegenerate) solutions for $n \geq 4$. We shall now solve them explicitly for $n = 2, 3$.

The case $n = 2$: In this case the roots are $\pm \alpha$ ($\alpha = e_1 - e_2$) and the structure constants on the right hand side of (10) vanish giving the one equation

$$\alpha \cdot (R^{\alpha} - R^{-\alpha}) = 0. \quad (11)$$

The vanishing of the structure constants means equations (9) and (11) decouple and we may solve (9) in terms of an arbitrary function f by

$$R^{\alpha\alpha} = -R^{-\alpha-\alpha} = f \quad \text{and} \quad R^{\alpha-\alpha} = -R^{-\alpha\alpha} = f + \frac{w'_\alpha}{w_\alpha}. \quad (12)$$

Similarly we may solve (11) in terms of arbitrary functions g_1, g_2 and h by

$$R^{i\alpha} = g_i \quad \text{and} \quad R^{i-\alpha} = g_i + h. \quad (13)$$

Finally $P^{ij} = \begin{pmatrix} \epsilon_1 & \epsilon_2 \\ \epsilon_1 & \epsilon_2 \end{pmatrix}$ satisfies $\alpha \cdot P^j = 0$. In matrix form this means the R -matrix looks like

$$\begin{pmatrix} \epsilon_1 & g_1 & 0 & f \\ g_1 + h & \epsilon_2 & f + \frac{w'_\alpha}{w_\alpha} & 0 \\ 0 & -f - \frac{w'_\alpha}{w_\alpha} & \epsilon_1 & g_2 \\ -f & 0 & g_2 + h & \epsilon_2 \end{pmatrix}. \quad (14)$$

By choosing $\epsilon_1 = \epsilon_2 = f = 0$ and $g_1 = g_2 = -h/2 = -w_\alpha$ we obtain an R -matrix in the form of Avan and Talon's [8, eq.(17)].

The case $n = 3$: Corresponding to any root $\alpha \in \Phi$ there are now precisely two roots $\gamma_{1,2}$ such that $\alpha + \gamma_{1,2} \in \Phi$. Indeed $\gamma_1 + \gamma_2 = -\alpha$. This peculiarity of $n = 3$ allows us to make the following ansatz for $R^{\alpha\alpha}$:

$$R^{\alpha\alpha} = -\frac{1}{2} \left(\frac{w'_\alpha}{w_\alpha} + \frac{w'_{\gamma_1}}{w_{\gamma_1}} + \frac{w_\alpha w_{\gamma_1}}{w_{\alpha+\gamma_1}} \right) = -\frac{1}{2} \left(\frac{w'_\alpha}{w_\alpha} + \frac{w'_{\gamma_2}}{w_{\gamma_2}} + \frac{w_\alpha w_{\gamma_2}}{w_{\alpha+\gamma_2}} \right). \quad (15)$$

The righthand equality between the two expressions involving the different roots $\gamma_{1,2}$ follows from the addition formula (2) together with the identity $z_\alpha = \frac{3}{2}\wp(u) + w_\alpha^2$ for $u \in \{\omega_1, \omega_2, -\omega_1 - \omega_2\}$. Thus $R^{\alpha\alpha}$ is well defined¹ and depends only on the choice of α . Further $R^{\alpha\alpha} + R^{-\alpha-\alpha} = 0$ and the remaining equations of (9) are solved upon setting

$$R^{-\alpha\alpha} = \frac{1}{2} \left(-\frac{w'_\alpha}{w_\alpha} + \frac{w'_{\gamma_1}}{w_{\gamma_1}} + \frac{w_\alpha w_{\gamma_1}}{w_{\alpha+\gamma_1}} \right) = \frac{1}{2} \left(-\frac{w'_\alpha}{w_\alpha} + \frac{w'_{\gamma_2}}{w_{\gamma_2}} + \frac{w_\alpha w_{\gamma_2}}{w_{\alpha+\gamma_2}} \right). \quad (16)$$

Our ansatz has the following two important features: if $\beta = \alpha + \gamma \in \Phi$ (for some γ), then $R^{\alpha\alpha} + R^{\beta\beta} = 0$ while $R^{-\alpha\alpha} + R^{\beta-\beta} = w_\alpha w_\beta / w_\gamma$. Note this means the first term on the righthand side of (10) will *always* vanish. This feature appears to hold for all of the R -matrix ansatz for Calogero-Moser related models we know of; it is usually achieved by setting $R^{\alpha\alpha} = 0$. Utilising these features means (10) simplifies to

$$\alpha \cdot \frac{R^{-\beta}}{w_{-\beta}} + \beta \cdot \frac{R^\alpha}{w_\alpha} = -c_{\alpha\gamma}^\beta. \quad (17)$$

This equation also arises (perhaps implicitly) in all of the R -matrix ansatz we know of. For gl_n a solution to the equation

$$\alpha \cdot A^{-\beta} + \beta \cdot A^\alpha = c_{\alpha\gamma}^\beta \quad (18)$$

¹ In the notation of [10] this choice corresponds to $\mathcal{A} = -\frac{3}{2}\wp(u)$.

is given by $A^{i\alpha} = |e_i \cdot \alpha|/2$ and so $R^{i\alpha} = -|e_i \cdot \alpha|w_\alpha/2$ yields a possible solution to (17). Actually (18) has a more general solution. Because the vector $(1, 1, 1)$ is orthogonal to every root then we may add a multiple of this to every solution of (18). Thus $R^{i\alpha} = -|e_i \cdot \alpha|w_\alpha/2 + f^\alpha$ is also a solution for arbitrary functions f^α (one for each root). Further $A^{i\alpha} = \alpha_i f^i$ will satisfy the homogeneous part of (18) for arbitrary functions f^i and so we obtain the solution

$$R^{i\alpha} = -|e_i \cdot \alpha|w_\alpha/2 + f^\alpha + \alpha_i f^i w_\alpha \quad (19)$$

in terms of the nine arbitrary functions f^α, f^i . A similar solution holds for arbitrary n . Observe that (17) contains 9 constraints on the 18 components $R^{i\alpha}$, six of them corresponding to three different roots which sum to zero and three coming from $\alpha = \beta$ for α a positive root. This accounts for the nine arbitrary functions in our solution.

Finally $P^{ij} = \begin{pmatrix} \epsilon_1^1 & \epsilon_2^2 & \epsilon_3^3 \\ \epsilon_1^1 & \epsilon_2^2 & \epsilon_3^3 \\ \epsilon_1^1 & \epsilon_2^2 & \epsilon_3^3 \end{pmatrix}$ satisfies $\alpha \cdot P^j = 0$. This means R -matrix takes the form given in Table 1, where for $\alpha = e_i - e_j$ we have set $w_\alpha \equiv w_{ij}$, $f^\alpha \equiv f^{ij}$ and $R^{\alpha\alpha} = \mathcal{R}_{ij}$. We note that as the elliptic potential degenerates, $\lim_{k \rightarrow 0} R^{\alpha\alpha} = 0$. Therefore the R -matrix obtained by setting $\epsilon_i = f^i = f^\alpha = 0$ yields the solution of Avan and Talon for the type I, II and III potentials.

To conclude. In this letter, we have presented general momentum independent R -matrices for the $n = 2$ and $n = 3$ Calgero-Moser systems with elliptic (type IV) potentials of Jacobi type, *i.e.*, $w(q) = -w(-q)$. These R -matrices reduce to the known solutions given by Avan and Talon for the potentials of the type I, II and III and are the first R -matrices known for these two particular models. The R -matrices found here have $R^{\alpha\alpha} \neq 0$ for the general elliptic potential. This feature is new: the R -matrices normally associated with the Calogero-Moser models have $R^{\alpha\alpha} = 0$. The nontrivial ansatz (15) makes full use of the $n = 3$ root space geometry and it is the particularly simple structure of the $n = 2, 3$ root spaces that allows a momentum independent solution.

T. S. acknowledges financial support from both the Daiwa Anglo-Japanese Foundation and Fujukai Foundation.

References

- [1] F. Calogero. Exactly solvable one-dimensional many-body problems. *Lett. Nuovo Cim.* **13** (1975) 411–416
- [2] J. Moser. Three integrable Hamiltonian systems connected with isospectral deformations. *Adv. Math.* **16** (1975) 1–23
- [3] M. A. Olshanetsky, A. M. Perelomov. Classical integrable finite dimensional systems related to Lie algebras. *Phys. Rep.* **71** (1981) 313–400

- [4] I. M. Krichever. Elliptic solutions of the Kadomtsev-Petviashvili equation and integrable systems of particles. *Func. Anal. Appl.* **14** (1980) 282–290
- [5] M. A. Semenov-Tian-Shansky. What is a classical r-matrix. *Funct. Anal. Appl.* **17** (1983) 259.
- [6] L. D. Faddeev, L. A. Takhtajan. *Hamiltonian methods in the theory of solitons*. Berlin, Heidelberg, New York: Springer 1987.
- [7] O. Babelon, C.-M. Viallet. Hamiltonian structures and Lax equations. *Phys. Lett.* **237** (1990) 411–416
- [8] J. Avan, M. Talon. Classical R -matrix structure for the Calogero model. *Phys. Lett.* **B303** (1993) 33–37
- [9] E. K. Sklyanin. Dynamical r-matrices for the elliptic Calogero-Moser Model. *LPTHE-93-42*; [hep-th/9308060](#)
- [10] H. W. Braden and T. Suzuki. R -matrices for Elliptic Calogero-Moser Models. *Edinburgh-/92-93/03*; [hep-th/9309033](#) To appear, *Letters in Mathematical Physics*.

This figure "fig1-1.png" is available in "png" format from:

<http://arXiv.org/ps/hep-th/9312031v1>